

# Benadryl Dosages

Product	Children's Benadryl Allergy Liquid	Children's Benadryl Allergy Liquid Dye-Free	Children's Benadryl-D Allergy & Sinus	Children's Benadryl Allergy Chewables	Children's Benadryl Allergy Fastmelt	Children's Benadryl Allergy & Cold Fastmelt
Symptoms	itchy, watery eyes runny nose itchy throat  sneezing	itchy, watery eyes runny nose itchy throat  sneezing	itchy, watery eyes runny nose itchy throat sinus pressure sneezing	itchy, watery eyes runny nose itchy throat  sneezing	itchy, watery eyes runny nose itchy throat  sneezing	itchy, watery eyes runny nose  sinus pressure sneezing cough

<b>DOSAGES</b>	Children 6 to under 12 years (> 50 lbs)	1-2 tsp every 4-6 hours as needed	1-2 tsp every 4-6 hours as needed	1-2 tsp every 4-6 hours as needed	1-2 tablets every 4-6 hours as needed	1-2 tablets every 4-6 hours as needed	1-2 tablets every 4-6 hours as needed
	Children 2 to 5 years (35 - 50 lbs)	3/4 tsp every 6 hours as needed	3/4 tsp every 6 hours as needed	3/4 tsp every 6 hours as needed	1 tablet every 6 hours as needed	1 tablet every 6 hours as needed	1 tablet every 6 hours as needed
	Children 6 mths to 2 yrs (20 - 35 lbs)	1/2 tsp every 6 hours as needed	1/2 tsp every 6 hours as needed	1/2 tsp every 6 hours as needed	not recommended	not recommended	not recommended
	Children 3-6 months (10 - 20 lbs)	1/4 tsp every 6 hours as needed	1/4 tsp every 6 hours as needed	1/4 tsp every 6 hours as needed	not recommended	not recommended	not recommended

**\*\* Do not exceed 4 doses in 24 hours for all products**

lbs=pounds

tsp=teaspoonful

mths=months

1 dropperful=0.8 ml

1 teaspoonful=5 ml